




Universidad
de Oviedo

SEARCHES FOR STANDARD MODEL HIGGS BOSON AT THE TEVATRON

Bárbara Álvarez González

Universidad de Oviedo

Prueba de Suficiencia Investigadora,
Octubre, 2007




CDF Top Event in 1995 on Computer Screen

Outline

- Tevatron.
- CDF Detector.
- Motivation of the Higgs Search at Tevatron.
- Matrix Element Analysis.
 - ◇ Application of the method to Single top search.
- Conclusions.

Tevatron


- The world's highest-energy particle accelerator.




- $p\bar{p}$ (proton-antiproton) collisions at 1.96 TeV center of mass energy.
- Collisions every 396 ns.
- Two collision points: **CDF** and **DØ**.
- Run I (1992 - 1996): Top quark was discovered
- Run II (2001 - nowadays): top quark and W mass, single top and ZZ production evidence and B_s mixing.
- Now, It is time for Higgs.

Tevatron Performance

TOTAL LUMI DELIVERED / STORED




DATA TAKING EFFICIENCY



- $\sim 3.2 \text{ fb}^{-1}$ delivered ($\sim 2.8 \text{ fb}^{-1}$ recorded).
- Highest instantaneous luminosity $\sim 3 \times 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$.
- $N_{events} = L \cdot \sigma$

CDF Detector (Collider Detector at Fermilab)

- General purpose particle detector with cylindrical symmetry.
- 3 subsystems: tracking (inside a 1.4 T solenoidal magnetic field), calorimetry and muons systems
- For Higgs physics the **full detector** is needed.



Motivation

- The Higgs boson is the last remaining Standard Model (SM) particle to be discovered.


- Direct limit for LEP2:

$$m_H > 114.4 \text{ GeV}/c^2 \text{ at } 95\% \text{ CL.}$$

- Considering the new W and top mass measurements


$$m_H < 144.4 \text{ GeV}/c^2 \text{ at } 95\% \text{ CL}$$

- Light Higgs is preferred.



Current Situation at Tevatron

- $D\bar{0}$ and CDF have established the search for the SM Higgs as one of their **highest priorities**.



- No single improvement by itself will reach the sensitivity needed. All channels studied by both experiments must be combined, and as much data as possible must be analyzed.

SM Higgs Physics: Production and Decay



- Low mass: $m_H < 135$ GeV

- ◆ The most relevant production mechanism is the associated production together a vector boson (W or Z).
- ◆ The highest branching ratio decay channel is to $H \rightarrow b\bar{b}$.
- ◆ The main backgrounds are $Wb\bar{b}$, $t\bar{t}$, single top, $Z\tau\tau$ and dibosons.

- High mass: $135 < m_H < 200$ GeV

- ◆ Production mechanism $gg \rightarrow H$
- ◆ The predominant decay channel is to $H \rightarrow WW$.
- ◆ The main backgrounds are dibosons, Drell-Yan, $t\bar{t}$ and single top.

0.8 - 0.2 pb for $gg \rightarrow H$
 0.2 - 0.03 pb for WH
 0.1 - 0.01 pb for ZH



Background Estimate

Top/EWK (WW/WZ/Z $\rightarrow\tau\tau$, ttbar, single t)



- MC normalized to theoretical cross-section

Non-W (QCD)

- Multijet events with semileptonic b -decays or mismeasured jets
- Fit low missing E_T data and extrapolate into signal region


W+HF jets (Wbb/Wcc/Wc)

- W+jets normalization from data and heavy flavor (HF) fractions from ALPGEN Monte Carlo, calibrated in generic multijet data



Mistags (W+2jets)

- Falsely tagged light quark or gluon jets
- Mistag probability parameterization obtained from inclusive jet data



Event Selection

- Event Selection:

- ◇ High p_T isolated lepton (e/μ):
 $p_T > 20$ GeV.


- ◇ High missing transverse energy:
 $E_T > 25$ GeV.

- ◇ Two jets with:


- ★ Transverse energy:
 $E_T > 20$ GeV.

- ★ Pseudorapidity ($\eta \equiv -\ln \tan(\frac{\theta}{2})$):
 $|\eta| < 2.8$

- ◇ At least one jet identified as b-jet.



Very challenging!!!




Sophisticated Tools Needed

- Signal is much smaller than background.

Signal hidden behind background uncertainty!

◇ Counting experiment are not possible \Rightarrow need sophisticated tools to isolate regions with high signal purity.



- We use a Matrix Element (ME) Technique to calculate event probabilities for the signal and the background (bkg) hypothesis.

Matrix Element Method

- Start from Fermi's Golden rule:

$$P_{evt} = \frac{d\sigma}{\sigma} = \frac{1}{\sigma} |M|^2 d\Phi$$

- But we need to consider:

- ◊ Parton distribution functions (PDFs): interactions are initiated by partons inside the p and \bar{p} .

- ◊ Neutrinos in the final state are not identified directly.

- ◊ The energy resolution of the detector can not be ignored.

- Sum over all possible particle variables (y) leading to the observed variables (x).

$$P_{evt} = \frac{1}{\sigma} \int d\sigma(y) dq_1 dq_2 f(x_1) f(x_2) W(y, x)$$

- $f(x_i)$ are the PDFs with $xi = \frac{E_{q_i}}{E_{beam}}$ and $W(y, x)$ is the transfer function (TF).
- The transfer function (TF) is the probability of measuring x when y was produced.

Transfer Function

- The transfer function is given by:

$$W(y, x) = \delta^3(p_l^y - p_l^x) \Pi_{i=1}^2 \delta^2(\Omega_i^y - \Omega_i^x) \Pi_{j=1}^2 W_{jet}(E_{parton_j}, E_{jet_j})$$

- " y " represents all final state particle momenta at particle level and " x " represents the measured momenta.
- p_l^y and p_l^x are the produced and measured lepton momenta.
- Ω_i^y and Ω_i^x are the produced quark and measured jet angles.
- E_{parton} and E_{jet} are the produced quark and measured jet energies.
- Because the lepton and jet angles are well measured:



$$W(y, x) = \Pi_{j=1}^2 W_{jet}(E_{parton_j}, E_{jet_j})$$

- The TF correct for jet energy, hadronization, detector effects,...

Application to the search of the Single Top

- s-channel single top has the same final state as $WH \rightarrow l\nu b\bar{b}$
- Benchmark for $WH \rightarrow l\nu b\bar{b}$

single top (s – channel)







- The complete application of the ME method to this final state has shown an evidence single top process.
- Transition from top physics to higgs physics.

Single Top Event Selection and Kinematic Variables

- Same event selection as $WH \rightarrow \nu b b \bar{b}$:


- ◇ One Lepton
 $p_T > 20 \text{ GeV}$,
 $|\eta_{e(m)}| < 2.0(1.0)$
- ◇ $\cancel{E}_T > 25 \text{ GeV}$.
- ◇ Two jets
 $E_T > 20 \text{ GeV}$,
 $|\eta| < 2.8$
- ◇ At least one jet identified as b-jet.
- ◇ This kinematic variables are the input variables for the event probabilities.



Event Probability Discriminant

- Define ratio of probabilities, obtained with the ME, as **Event Probability Discriminant (EPD)**:


$$EPD = \frac{b P_{single\ top}}{b P_{single\ top} + b P_{Wbb} + (1-b) P_{Wcc} + (1-b) P_{Wcj}}$$



Binned Likelihood Fit

- Likelihood function:



$$L(\beta_1, \dots, \beta_5; \delta_1, \dots, \delta_{10}) = \prod_{k=1}^B \frac{e^{-\mu_k} \mu_k^{n_k}}{n_k!} \cdot \prod_{j=2}^5 G(\beta_j | 1, \Delta_j) \cdot \prod_{i=1}^{10} G(\delta_i | 0, 1)$$



Hypothesis Test

- We use the CLs/CLb Method developed at LEP.
- Define Likelihood ratio test statistic \Rightarrow
- Most sensitive bins:

$$Q = \frac{L(data|s+b)}{L(data|b)}$$



Expected sensitivity: 0.10% (3.1 σ)

Observed sensitivity: 0.07% (3.2 σ)

$$WH \rightarrow l\nu b\bar{b}$$

- Application of the ME method and improvements for WH search.
- Improvements to Event Selection:
 - ◇ Add new lepton categories.
 - ◇ Reduce the backgrounds by improving the identification of b-jets, using new btagging algorithms.
- Improvements to the ME method:
 - ◇ Add information to the TFs.
 - ★ Different jet cone sizes for better energy resolution.
- Study of the mass resolution of the $b\bar{b}$ system.
- Study the systematic uncertainties.
- Add as much data as possible.
- Extract the maximal sensitivity from the data.



Improvements

- New lepton categories:


- ◇ **Electrons**: CEM and PHX.
- ◇ **Muons**: CMUP, CMX and BMU.

- Different cone size jets:

- ◇ Jets of conesize 0.4
- ◇ Jets of conesize 0.7



Parton Energy - Jet Energy




Conclusions

- Advanced analysis tools essential to establish small signals buried underneath large backgrounds.
- No single golden kinematic variable. Attempt to include all available kinematic information by using Matrix Element approach.
- Recent **evidence** for electroweak single top quark production at the Tevatron established by CDF and DØ experiment.
- Then, the Matrix Element Method has been validated by this evidence of the single top having the same final state as $WH \rightarrow l\nu b\bar{b}$

Conclusions

- Important milestone along the way to the Higgs!

- ◇ Most of the tools needed for the WH full analysis are ready.
- ◇ But several improvements, with respect to the single top analysis, are being studied.



Back Up



Fermi's Golden Rule

- And the Fermi's golden rule tell us the differential cross-section for a given scattering process:


$$d\sigma = \frac{(2\pi)^4 |M|^2}{4\sqrt{(q_1 \cdot q_2) - m_{q_1}^2 m_{q_2}^2}} d\Phi_n(q_1 + q_2; p_1, \dots, p_n)$$

- ◇ $|M|$ is the Lorentz invariant matrix element.
- ◇ q_1, q_2 and m_{q_1}, m_{q_2} are the four momenta and masses of the incident particles.
- ◇ $d\Phi_n$ is the n-body phase space.

Background Feynman Diagrams and Cross Sections



t \bar{t}




Process	Cross Section (pb)
<i>WH</i>	<i>1 - 0.01</i>
Single Top W^* (s-channel)	0.88 ± 0.11
Single Top $W - g$ (t-channel)	1.98 ± 0.25
ZZ	1.58 ± 0.02
WZ	3.96 ± 0.06
WW	13.25 ± 0.25
$t\bar{t}$	8.9 ± 1.0
$Z \rightarrow \tau^+ \tau^-$	254.3 ± 5.4

WH Event Probability Discriminant

- Definition of the discriminant: Event Probability Discriminant.

$$EPD = \frac{P_{evt}(signal)}{P_{evt}(signal) + P_{evt}(bkg)}$$




◇ The blue lines are the background and the red ones are the signal.

◇ The dotted line is the new TF and the solid line is the old TF.

◇ For the bottom plot, the red line is the new TF and the blue line is the old TF.

Transfer Function for jet energies



- Mapping between parton and jet energies:
 $\delta_E = E_{parton} - E_{jet}$
- We parameterize the δ_E distribution as a sum of two Gaussian functions:
 - ◊ One to account the peak.
 - ◊ The other to account the asymmetric tail.



- Transfer Function (TF):

$$W_{jet}(E_{parton}, E_{jet}) = \frac{1}{\sqrt{2\pi}(p_2 + p_3 p_5)} \left(\exp \frac{-(\delta_E - p_1)^2}{2p_2^2} + p_3 \cdot \exp \frac{-(\delta_E - p_4)^2}{2p_5^2} \right)$$

Where $p_i = a_i + b_i E_{parton}$. A total of 10 parameters ($a_1, b_1, \dots, a_5, b_5$) are required to specify $W_{jet}(E_{parton}, E_{jet})$.


Transfer Function Improvements

- We are adding more information to the mapping between partons and jets:
 - ◇ Jets of 0.7 conesize.
 - ◇ Different η regions.
 - ◇ SLT
 - ◇ SumE studies.
 - ◇ btagging variables: JetProb and SecVtx.




Transfer Function in parton energy bins

- Jets of 0.4 conesize, $\Delta R(\text{jet}, \text{parton}) < 0.4$ and $|\eta_{\text{jet}}| < 0.9$:
- Different parton energies (20,30,50,70,100,120,150,170,200,250)GeV.
- Parton Energy distribution ($n(E_{\text{parton}})$) fitted with two gaussians and one exponential.




TF in 2-D



E_{parton} distribution

Test the Transfer Function


- The performance of the parameterization is shown in the figure.
- The δ_E distribution is compared with the prediction from the transfer function.
- ◊ The histogram corresponds with the δ_E distribution.
- ◊ The solid line is the convolution of the transfer function with the parton energy spectrum.



$$H(E_{parton}, E_{jet}) = \int_{E_{parton}^1}^{E_{parton}^2} n(E_{parton}) \cdot W(E_{parton} - E_{jet}, E_{parton}) dE_{parton}$$


Calorimetry

- Scintillator based calorimeter: Scintillating detectors after absorbing material.
- As the particle goes through the absorbing material, it forms a shower of secondary particles.
- The scintillators convert the energy into light which can be observed by light-sensitive detectors.
- The amount of light observed measures the energy of the particle.



Particle identification with calorimeter

- In calorimeters different particles travel different distances before being absorbed.
- Photons and electrons lose energy very quickly and stop in the electromagnetic calorimeter.
 - ◊ Photons don't leave signal in the tracking detector.
- Muons can pass through many cm of material before losing their energy.
- Jets from quarks reach the hadron calorimeter and form hadronic showers.
- The distance a particle travels in a calorimeter is used to identify the particle.



CDF calorimeter

- Electromagnetic (EM) and Hadronic (HADRON) calorimeters :

- ◇ Lead for EM
- ◇ Steel for HADRON

- Coverage:

- ◇ $|\eta| < 3.6$; $\eta \equiv -\ln \tan(\frac{\theta}{2})$


- ◇ $\phi < 2\pi$

- Regions:

- ◇ Central: $|\eta| < 1.1$


- ◇ Plug: $1.1 < |\eta| < 2.1$

- ◇ Forward: $2.1 < |\eta| < 3.6$



Taggers

- **Soft Lepton** Tagger: looks for an energetic lepton inside a jet.
- Identify a b-jet with different btagging algorithms: Secvtx, JetProb and combined.



- $b \rightarrow \ell \nu c$ (BR $\sim 20\%$)
- $b \rightarrow c \rightarrow \ell \nu s$ (BR $\sim 20\%$)

